



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Mircera Page: 1 of 2

Effective Date: 3/4/2024 Last Review Date: 01/12/2024

Applies to: Illinois Florida Michigan
 New Jersey Maryland Florida Kids
 Pennsylvania Kids Virginia Texas

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Mircera under the patient's prescription drug benefit.

Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

Mircera is indicated for the treatment of anemia associated with chronic kidney disease (CKD) in:

- Adult patients on dialysis and adult patients not on dialysis.
- Pediatric patients 5 to 17 years of age on hemodialysis who are converting from another erythropoiesis-stimulating agent (ESA) after their hemoglobin level was stabilized with an ESA.

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Mircera

Policy/Guideline:

Criteria for Initial Approval:

Note: Requirements regarding hemoglobin level exclude values due to recent transfusion. All members must be assessed for iron deficiency anemia and have adequate iron stores (defined as a serum transferrin saturation [TSAT] level greater than or equal to 20% within the prior 3 months) or are receiving iron therapy before starting Mircera. Members may not use Mircera concomitantly with other erythropoiesis stimulating agents.

Anemia Due to Chronic Kidney Disease (CKD)

Authorization of 12 weeks may be granted for the treatment of anemia due to chronic kidney disease in members with pretreatment hemoglobin < 10 g/dL and patient is unable to take Epogen and Procrit for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication.

Continuation of Therapy:

Note: Requirements regarding current hemoglobin level exclude values due to recent transfusion. All members must be assessed for iron deficiency anemia and have adequate



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Mircera

Page: 2 of 2

Effective Date: 3/4/2024

Last Review Date: 01/12/2024

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Michigan
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Texas

iron stores (defined as a serum transferrin saturation [TSAT] level greater than or equal to 20% with the prior 3 months) or are receiving iron therapy before continuation of treatment with Mircera.

Members may not use Mircera concomitantly with other erythropoiesis stimulating agents.

Anemia Due to Chronic Kidney Disease (CKD)

1. Authorization of 12 weeks may be granted for continued treatment of anemia due to chronic kidney disease in members with current hemoglobin < 12 g/dL and the member has shown a response to therapy with a rise in hemoglobin of ≥ 1 g/dL after at least 12 weeks of ESA therapy.
2. Authorization of up to 12 weeks may be granted for continued treatment of anemia due to chronic kidney disease in members who have not completed 12 weeks of ESA therapy.

Approval Duration and Quantity Restrictions:

Approval: 12 weeks

References:

1. Mircera [package insert]. St. Gallen, Switzerland: Vifor (International) Inc.; March 2023.
2. Kidney Disease: Improving Global Outcomes (KDIGO) Anemia Work Group. KDIGO Clinical Practice Guideline for Anemia in Chronic Kidney Disease. *Kidney Int.* 2012;Suppl 2:279-335.