



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Otezla Page: 1 of 6

Effective Date: 2/1/2024 Last Review Date: 11/2023

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Otezla under the patient's prescription drug benefit.

Description:

FDA-Approved Indications

- A. Adult patients with plaque psoriasis (PsO) who are candidates for phototherapy or systemic therapy (Reference the Biological Response Modifiers (BRMs) in the Treatment of Plaque Psoriasis NJ Protocol)
- A. Adults with active psoriatic arthritis
- B. Adult patients with oral ulcers associated with Behcet's disease

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Preferred: Otezla

Policy/Guideline:

Documentation:

- A. **Psoriatic arthritis (PsA)**
 - 1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
 - 2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.
- B. **Behcet's disease** (initial requests only): Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy (if applicable).

Prescriber Specialty:

This medication must be prescribed by or in consultation with one of the following:

- A. Psoriatic arthritis: rheumatologist or dermatologist
- B. Bechet's disease: rheumatologist

Criteria for Initial Approval:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Otezla

Page: 2 of 6

Effective Date: 2/1/2024

Last Review Date: 11/2023

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

A. Psoriatic arthritis (PsA)

1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug indicated for active psoriatic arthritis.
2. Authorization of 12 months may be granted for adult members for treatment of active psoriatic arthritis when one of the following criteria is met:
 - i. Member has had an inadequate response to methotrexate, leflunomide, or another conventional synthetic drug (e.g., sulfasalazine) administered at an adequate dose and duration.
 - ii. Member has an intolerance or contraindication to methotrexate or leflunomide (see Appendix B), or another conventional synthetic drug (e.g., sulfasalazine).
 - iii. Member has enthesitis.

B. Behcet's disease

1. Authorization of 12 months may be granted for adult members who have previously received a biologic indicated for treatment of Behcet's disease.
2. Authorization of 12 months may be granted for adult members for treatment of oral ulcers associated with Behcet's disease when the member has had an inadequate response to at least one nonbiologic medication for Behcet's disease (e.g., colchicine, systemic glucocorticoids, azathioprine).

Continuation of Therapy:

A. Psoriatic arthritis (PsA)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for psoriatic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Axial disease
6. Skin and/or nail involvement

B. Behcet's disease



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Otezla Page: 3 of 6

Effective Date: 2/1/2024 Last Review Date: 11/2023

Applies to: Illinois Florida Florida Kids
 New Jersey Maryland Michigan
 Pennsylvania Kids Virginia Kentucky PRMD

Authorization of 12 months may be granted for all adult members (including new members) who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition.

Other Criteria:

Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug.

Appendix A: Table. Relative potency of select topical corticosteroid products

Potency	Drug	Dosage form	Strength
I. Super-high potency (group 1)	Augmented betamethasone dipropionate	Ointment, Lotion, Gel	0.05%
	Clobetasol propionate	Cream, Gel, Ointment, Solution, Cream (emollient), Lotion, Shampoo, Foam, Spray	0.05%
	Fluocinonide	Cream	0.1%
	Flurandrenolide	Tape	4 mcg/cm ²
	Halobetasol propionate	Cream, Lotion, Ointment, Foam	0.05%
II. High potency (group 2)	Amcinonide	Ointment	0.1%
	Augmented betamethasone dipropionate	Cream	0.05%
	Betamethasone dipropionate	Ointment	0.05%
	Clobetasol propionate	Cream	0.025%
	Desoximetasone	Cream, Ointment, Spray	0.25%
		Gel	0.05%
	Diflorasone diacetate	Ointment, Cream (emollient)	0.05%
	Fluocinonide	Cream, Ointment, Gel, Solution	0.05%
	Halcinonide	Cream, Ointment	0.1%
Halobetasol propionate	Lotion	0.01%	
III. High potency (group 3)	Amcinonide	Cream, Lotion	0.1%
	Betamethasone dipropionate	Cream, hydrophilic emollient	0.05%
	Betamethasone valerate	Ointment	0.1%
		Foam	0.12%
	Desoximetasone	Cream, Ointment	0.05%
	Diflorasone diacetate	Cream	0.05%
Fluocinonide	Cream, aqueous emollient	0.05%	



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Otezla Page: 4 of 6

Effective Date: 2/1/2024 Last Review Date: 11/2023

Applies to: Illinois Florida Florida Kids
 New Jersey Maryland Michigan
 Pennsylvania Kids Virginia Kentucky PRMD

Potency	Drug	Dosage form	Strength
	Fluticasone propionate	Ointment	0.005%
	Mometasone furoate	Ointment	0.1%
	Triamcinolone acetonide	Cream, Ointment	0.5%
IV. Medium potency (group 4)	Betamethasone dipropionate	Spray	0.05%
	Clocortolone pivalate	Cream	0.1%
	Fluocinolone acetonide	Ointment	0.025%
	Flurandrenolide	Ointment	0.05%
	Hydrocortisone valerate	Ointment	0.2%
	Mometasone furoate	Cream, Lotion, Solution	0.1%
	Triamcinolone acetonide	Cream	0.1%
		Ointment	0.05% and 0.1%
V. Lower-mid potency (group 5)	Betamethasone dipropionate	Lotion	0.05%
		Cream	0.1%
	Desonide	Ointment, Gel	0.05%
	Fluocinolone acetonide	Cream	0.025%
	Flurandrenolide	Cream, Lotion	0.05%
	Fluticasone propionate	Cream, Lotion	0.05%
	Hydrocortisone butyrate	Cream, Lotion, Ointment, Solution	0.1%
	Hydrocortisone probutate	Cream	0.1%
	Hydrocortisone valerate	Cream	0.2%
	Prednicarbate	Cream (emollient), Ointment	0.1%
	Triamcinolone acetonide	Lotion	0.1%
Ointment		0.025%	
VI. Low potency (group 6)	Alclometasone dipropionate	Cream, Ointment	0.05%
	Betamethasone valerate	Lotion	0.1%
	Desonide	Cream, Lotion, Foam	0.05%
	Fluocinolone acetonide	Cream, Solution, Shampoo, Oil	0.01%
	Triamcinolone acetonide	Cream, lotion	0.025%
Hydrocortisone (base, greater than or equal to 2%)	Cream, Ointment, Solution	2.5%	
	Lotion	2%	



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Otezla Page: 5 of 6

Effective Date: 2/1/2024 Last Review Date: 11/2023

Applies to: Illinois Florida Florida Kids
 New Jersey Maryland Michigan
 Pennsylvania Kids Virginia Kentucky PRMD

Potency	Drug	Dosage form	Strength
VII. Least potent (group 7)	Hydrocortisone (base, less than 2%)	Cream, Ointment, Gel, Lotion, Spray, Solution	1%
		Cream, Ointment	0.5%
	Hydrocortisone acetate	Cream	2.5%
		Lotion	2%
		Cream	1%

Appendix B: Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine, Acitretin, or Leflunomide

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
2. Drug interaction
3. Risk of treatment-related toxicity
4. Pregnancy or currently planning pregnancy
5. Breastfeeding
6. Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
7. Hypersensitivity
8. History of intolerance or adverse event

Approval Duration and Quantity Restrictions:

Approval:

Initial and Renewal Approval: 12 months

Quantity Level Limit:

Otezla (apremilast) starter pack: 1 pack (55 tablets) per 28 days

Otezla (apremilast) 30 mg tablets: 60 tablets per 30 days

FDA-Recommended Dosing:

Day 1 to day 5 dosage titration schedule

- Day 1: 10 mg in morning
- Day 2: 10 mg in morning and 10 mg in evening
- Day 3: 10 mg in morning and 20 mg in evening
- Day 4: 20 mg in morning and 20 mg in evening
- Day 5: 20 mg in morning and 30 mg in evening



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Otezla Page: 6 of 6

Effective Date: 2/1/2024 Last Review Date: 11/2023

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input checked="" type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Day 6 and thereafter

- 30 mg in morning and 30 mg in evening

References:

1. Otezla [package insert]. Thousand Oaks, CA: Amgen Inc.; December 2021.
2. Coates LC, Kavanaugh A, Mease PJ, et al. Group for research and assessment of psoriasis and psoriatic arthritis 2015 treatment recommendation for psoriatic arthritis. *Arthritis Rheumatol.* 2016 May;68(5):1060-71.
3. Menter A, Gelfand JM, Connor C, et al. Joint AAD-NPF guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol.* 2020;82(6):1445-1486.
4. [Gossec L](#), Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. *Ann Rheum Dis.* 2020;79(6):700-712.
5. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheum.* 2018;71:5-32.
6. Hatemi G, Christensen R, Bodaghi, et al. 2018 update of the EULAR recommendations for the management of Behcet's syndrome. *Ann Rheum Dis.* 2018.; 77: 808-818.
7. Stein Gold L, Papp K, Pariser D, et al. Efficacy and safety of apremilast in patients with mild-to-moderate plaque psoriasis: Results of a phase 3, multicenter, randomized, double-blind, placebo-controlled trial. *J Am Acad Dermatol.* 2022;86(1):77-85. doi:10.1016/j.jaad.2021.07.040.
8. Elmets CA, Korman NJ, Prater EF, et al. Joint AAD-NPF Guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures. *J Am Acad Dermatol.* 2021;84(2):P432-470.
9. Topical Corticosteroids. *Drug Facts and Comparisons.* Facts & Comparisons [database online]. St. Louis, MO: Wolters Kluwer Health Inc; December 1, 2021. Accessed January 11, 2023.
10. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. *Nat Rev Rheumatol.* 2022;18(8):465-479.