



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Letairis

Page: 1 of 4

Effective Date: 11/1/2024

Last Review Date: 10/2024

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Letairis under the patient's prescription drug benefit.

Description:

FDA-Approved Indication

Indicated for the treatment of pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1):

- A. To improve exercise ability and delay clinical worsening
- B. In combination with tadalafil to reduce the risks of disease progression and hospitalization for worsening PAH, and to improve exercise ability.

Studies establishing effectiveness included trials predominantly in patients with WHO Functional Class II-III symptoms and etiologies of idiopathic or heritable PAH or PAH associated with connective tissue diseases.

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Letairis
ambrisentan

Policy/Guideline:

Prescriber Specialty:

This medication must be prescribed by or in consultation with a pulmonologist or cardiologist.

Criteria for Initial Approval:

Pulmonary Arterial Hypertension (PAH)

Authorization of 12 months may be granted for treatment of PAH when ALL of the following criteria are met:

- A. Member has PAH defined as WHO Group 1 class of pulmonary hypertension (refer to Appendix).
- B. PAH was confirmed by either criterion (1) or criterion (2) below:
 - 1. Pretreatment right heart catheterization with all of the following results:
 - i. mPAP > 20 mmHg



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Letairis

Page: 2 of 4

Effective Date: 11/1/2024

Last Review Date: 10/2024

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- ii. PCWP \leq 15 mmHg
- iii. Pulmonary vascular resistance (PVR) \geq 3 Wood units in adult patients or pulmonary vascular resistance index (PVRI) \geq 3 Wood units x m² in pediatric patients
- 2. For infants less than one year of age, PAH was confirmed by Doppler echocardiogram if right heart catheterization cannot be performed.
- C. Requests for generic Letairis (ambrisentan): Patient is unable to take the preferred brand Letairis for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication.

Criteria for Continuation of Therapy:

Authorization of 12 months may be granted for members with an indication listed in criteria for initial approval who are currently receiving the requested medication through a paid pharmacy or medical benefit, and who are experiencing benefit from therapy as evidenced by disease stability or disease improvement.

Appendix:

WHO Classification of Pulmonary Hypertension

1 PAH

- 1.1 Idiopathic (PAH)
- 1.2 Heritable PAH
- 1.3 Drug- and toxin-induced PAH
- 1.4. PAH associated with:
 - 1.4.1 Connective tissue diseases
 - 1.4.2 HIV infection
 - 1.4.3 Portal hypertension
 - 1.4.4 Congenital heart diseases
 - 1.4.5 Schistosomiasis
- 1.5 PAH long-term responders to calcium channel blockers
- 1.6 PAH with overt features of venous/capillaries (PVOD/PCH) involvement
- 1.7 Persistent PH of the newborn syndrome

2 PH due to left heart disease

- 2.1 PH due to heart failure with preserved LVEF
- 2.2 PH due to heart failure with reduced LVEF
- 2.3 Valvular heart disease
- 2.4 Congenital/acquired cardiovascular conditions leading to post-capillary PH



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Letairis Page: 3 of 4

Effective Date: 11/1/2024 Last Review Date: 10/2024

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

3 PH due to lung diseases and/or hypoxia

- 3.1 Obstructive lung disease
- 3.2 Restrictive lung disease
- 3.3 Other lung disease with mixed restrictive/obstructive pattern
- 3.4 Hypoxia without lung disease
- 3.5 Developmental lung disorders

4 PH due to pulmonary artery obstruction

- 4.1 Chronic thromboembolic PH
- 4.2 Other pulmonary artery obstructions
 - 4.2.1 Sarcoma (high or intermediate grade) or angiosarcoma
 - 4.2.2 Other malignant tumors
 - Renal carcinoma
 - Uterine carcinoma
 - Germ cell tumours of the testis
 - Other tumours
 - 4.2.3 Non-malignant tumours
 - Uterine leiomyoma
 - 4.2.4 Arteritis without connective tissue disease
 - 4.2.5 Congenital pulmonary artery stenosis
 - 4.2.6 Parasites
 - Hydatidosis

5 PH with unclear and/or multifactorial mechanisms

- 5.1 Hematologic disorders: Chronic hemolytic anemia, myeloproliferative disorders
- 5.2 Systemic and metabolic disorders: Pulmonary Langerhans cell histiocytosis, Gaucher disease, glycogen storage disease, neurofibromatosis, sarcoidosis
- 5.3 Others: chronic renal failure with or without hemodialysis, fibrosing mediastinitis
- 5.4 Complex congenital heart disease

Approval Duration and Quantity Restrictions:

Approval: 12 months

Quantity Level Limit:

- Letairis 5 mg tablets: 30 per 30 days
- Letairis 10 mg tablets: 30 per 30 days

References:

1. Letairis [package insert]. Foster City, CA: Gilead Sciences, Inc.; August 2019.
2. Ambrisentan [package insert]. Cranbury, NJ: Sun Pharmaceutical Industries, Inc.; January 2024.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Letairis Page: 4 of 4

Effective Date: 11/1/2024 Last Review Date: 10/2024

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

3. McLaughlin VV, Archer SL, Badesch DB, et al. ACCF/AHA 2009 expert consensus document on pulmonary hypertension a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association developed in collaboration with the American College of Chest Physicians, American Thoracic Society, Inc., and the Pulmonary Hypertension Association. *J Am Coll Cardiol.* 2009;53(17):1573-1619.
4. Badesch DB, Champion HC, Gomez-Sanchez MA, et al. Diagnosis and assessment of pulmonary arterial hypertension. *J Am Coll Cardiol.* 2009;54:S55-S66.
5. Taichman DB, Ornelas J, Chung L, et al. Pharmacologic therapy for pulmonary arterial hypertension in adults: CHEST guideline and expert panel report. *Chest.* 2014;146(2):449-475.
6. Abman SH, Hansmann G, Archer SL, et al. Pediatric pulmonary hypertension: guidelines from the American Heart Association and American Thoracic Society. *Circulation.* 2015;132(21):2037-99.
7. Klinger JR, Elliott CG, Levine DJ, et al. Therapy for pulmonary arterial hypertension in adults: update of the CHEST guideline and expert panel report. *Chest.* 2019;155(3):565-586.
8. Galie N, McLaughlin VV, Rubin LJ, Simonneau G. An overview of the 6th World Symposium on Pulmonary Hypertension. *Eur Respir J.* 2019;53(1):1802148. doi: 10.1183/13993003.02148-2018
9. Simonneau G, Montani D, Celermajer DS, et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J.* 2019;53(1):1801913. doi:10.1183/13993003.01913-2018